

Memorial EMS  
Decatur Memorial EMS  
Springfield Memorial EMS

## **OB/GYN PROTOCOLS**



Memorial EMS  
Decatur Memorial EMS  
Springfield Memorial EMS

Childbirth Protocol.....	17.A.1
Obstetrical Complications Protocol .....	17.B.1
Abnormal Delivery Protocol.....	17.C.1



## Childbirth Protocol

Childbirth is a natural process. EMS providers called to a woman in labor should determine whether there is enough time to transport the expecting mother to the hospital or if delivery is imminent. If childbirth appears imminent, immediately prepare to assist with the delivery.

### EMR Care

Care should be focused on assessing the situation, initiating routine patient care and preparing for or providing patient transport. Special attention should be given to the privacy of the mother and concerns of immediate family members should be addressed.

1. Render initial care in accordance with the *Routine Patient Care Protocol*.
2. **Oxygen:** If respiratory distress is noted, 15 LPM via NRM or if unable to tolerate the mask, 6 LPM via nasal cannula.
  - a) If no obvious respiratory distress is noted, apply a pulse ox. If  $\geq 94\%$  and no signs/ symptoms of respiratory distress, no Oxygen is required. If  $\leq 94\%$  apply nasal cannula at 2-6 LPM or 15 LPM via NRM as needed to raise pulse ox to  $\geq 94\%$ .
3. Obtain a history on the patient including:
  - a) Gravida (# of pregnancies) and Para (# of live births)
  - b) Expected delivery date
  - c) Length of previous labor and if complications of previous pregnancies
  - d) Onset of contractions
  - e) Prenatal care (if any)
4. Allow the expectant mother to remain in a position that is most comfortable.
  - a) If patient is supine, one provider should be dedicated to provide manual left uterine displacement, moving uterus to patients left side.
5. If delivery is not imminent, transport the patient on her left side.
6. Determine if there is adequate time to transport:
  - a) Assess the nature, extent and time of contractions.
  - b) Assess the patient for high-risk factors.
  - c) Assess the status of the membranes and any discharge.
  - d) Assess for pushing with contractions.
  - e) Take into consideration the length of previous labor.
7. If delivery is imminent:
  - a) Consider requesting additional ambulance.
  - b) **DO NOT ATTEMPT TO RESTRAIN OR DELAY DELIVERY**
  - c) Position the mother supine on a flat surface if possible.
  - d) Use full PPE – gloves, gown & goggles.

## Childbirth Protocol

### EMR Care (Continued)

8. Prepare for delivery:
  - a) Control delivery of the head so that it does not emerge too quickly. Support the infant's head as it emerges and protect the perineum with gentle hand pressure.
  - b) Assess for nuchal cord and, if present, gently remove the cord from around the newborn's neck.
  - c) Suction the mouth, then nose of the newborn with a bulb syringe as soon as the head is delivered.
  - d) As the shoulders emerge, guide the head & neck downward to deliver the anterior shoulder. Support and lift the head & neck slightly to deliver the posterior shoulder.
  - e) Ensure a firm hold on the baby as the rest of the newborn's body delivers.
  - f) Keep the newborn level with the mother's vagina until the cord stops pulsating and is double clamped.

### EMT Care

1. EMT Care includes all components of *EMR Care*.
2. **Contact Medical Control** as soon as possible.

### A-EMT/ EMT-I

1. A-EMT/ EMT-I includes all components of *EMT Care*.
2. **IV Fluid Therapy:** IVF 500mL fluid boluses to maintain a systolic BP of at least 100mmHg.
  - a) IV placement should be higher than the level of the diaphragm for medication administration.
  - b) While not a preferred location, if IO access is required, proximal humerus should be utilized (trauma, arrest, and/or hemorrhage situations).

### Paramedic Care

1. Paramedic Care includes all components of *A-EMT/ EMT-I Care*.

## Childbirth Protocol

EMR, **EMT**, **A-EMT/ EMT-I**, **PARAMEDIC CARE**

### Infant Post Partum Care

1. Begin the *Emergency Childbirth Record*.
2. Dry the newborn and wrap in a warm blanket, keeping the baby at the level of the mother's vagina until the cord is clamped and cut.
3. Continue to suction the nose and mouth. Spontaneous respirations should begin within *15 seconds*.
  - If spontaneous respirations are not present, begin artificial ventilations with **BVM**: 100% O<sub>2</sub> at 30-40 bpm.
  - If no brachial pulse is present **OR** the pulse is less than 100 bpm, begin **CPR**.
4. After the umbilical cord stops pulsating, clamp the cord at 3" & at 4" from the newborn's abdomen and cut between the clamps with the sterile scalpel found in the OB kit.
5. Assess the cord for bleeding and note the number of vessels present.
6. Obtain an APGAR score at **1 minute** and again at **5 minutes** after delivery.
7. Place ID tags on the mother and infant with the following information:
  - Name of the mother
  - Sex of the infant
  - Date and time of delivery
8. **DO NOT** separate the mother and infant unless both have ID tags.

### Post Partum Care of the Mother

1. The placenta should deliver within 5-20 minutes. Collect the placenta in a plastic bag and bring it to the hospital with the mother. **DO NOT** pull on the cord to facilitate delivery of the placenta.
2. Do not delay transport for delivery of the placenta.
3. If the perineum is torn and bleeding, apply direct pressure with a 5x9 dressing or trauma dressing and have the patient bring her legs together.

## Childbirth Protocol

Be alert, excessive **maternal bleeding (most commonly uterine atony)** is a **life-threatening condition** that requires aggressive treatment. Communication to the receiving facility and providing focused care for the mother will be required for successful resuscitation.

### EMR Care

1. Massage the fundus vigorously until firm.
  - a) *To massage the fundus, place one hand with fingers fully extended just above the mother's pubic bone and use the other hand to press down into the abdomen and gently massage the uterus approximately 3 to 5 minutes until it becomes firm.*
2. Keep the mother warm, baby to breast when possible.
3. **Oxygen:** Titrate O2 admin to maintain  $SpO_2 \geq 94\%$ .

### EMT Care

1. EMT Care includes all components of *EMR Care*.
2. Apply **Waveform Capnography**
3. **Contact Medical Control** as soon as possible.

### A-EMT/ EMT-I

1. A-EMT/ EMT-I includes all components of *EMT Care*.
2. Obtain **IV access**.
3. **IV Fluid Therapy:** Continue fluid bolus to maintain  $BP \geq 100\text{mmHg}$ .

### Paramedic Care

1. Paramedic Care includes all components of *A-EMT/ EMT-I Care*.
2. **Tranexamic Acid (TXA):** 1 gm in 100 mL D5W over 10 minutes IV infusion

**Inclusion criteria** for the administration of Tranexamic Acid:

- a) Patient age of 18 years or older.
- b) Ongoing hemorrhage with estimated blood loss  $> 500\text{mL}$ 
  - a. Sustained tachycardia  $HR > 120$ .
  - b. Sustained hypotension  $SBP < 90\text{mmHg}$ .
- c) Signs of peripheral vasoconstriction
  - a. Cool, pale skin.
  - b. Delayed cap refill.
- d) **Delivery occurred  $\leq 3$  hours prior to TXA administration.**

**Absolute exclusion criteria:**

- a) Time of delivery  $> 3$  hours or unknown.
- b) Patients who antifibrinolytic therapy is contraindicated.
  - a. A history of DVT/PE or procoagulant disorder (i.e. protein c, protein s or Antithrombin III disorder)
- c) Age  $< 18$

### Critical Thinking Elements

- TXA should never be administered at a "wide open" rate.
- Hypotension has been observed when TXA is administered too rapidly.
- Use with caution in patients with a history of DVT, PE, known clotting disorders, and severe renal failure.

## Childbirth Protocol

### High-Risk Pregnancy Factors

- Lack of prenatal care
- Drug abuse
- Teenage pregnancy
- Diabetes
- Hypertension
- Cardiac disease
- Previous breech or C-section delivery
- Pre-eclampsia / Toxemia / Eclampsia
- Twins / Multiple birth pregnancy

### Pre-Term OB

### Transporting Units

Destination decisions must be informed decisions based on local and regional destination capabilities, time since onset and transportation distances.

1. Patients meeting *Patient Destination Criteria (6.G.1)* should be transported to a Level III Perinatal facility if possible.
  - a. If unable to safely transport directly, include this in PCR documentation.
  - b. If the patient refuses, include documentation of the informed refusal in communication with **Medical Control** and in the PCR.

### Documentation Requirements

1. Completed *Emergency Childbirth Record* and attach to the PCR. (linked [here](#))
2. Document the date, time, and place of delivery.
3. Presence or absence of a nuchal cord.
  - If nuchal cord is present, document how many times the cord was wrapped around the baby's neck.
4. Appearance of the amniotic fluid.
5. Time the placenta was delivered and its condition.
6. APGAR score at **1 minute** and **5 minutes**.
7. Any resuscitation / treatment rendered and newborn response to treatment.

### Critical Thinking Elements

- Lower than normal blood pressure and higher than usual heart rate are normal vital sign changes with pregnancy.
- Signs & symptoms of shock in the pregnant patient include a systolic BP less than 90mmHg, lightheadedness and ALOC.
- Average labor lasts 8-12 hours but can be as short as 5 minutes.
- The desire to push during contractions is an indicator that delivery is imminent.
- Be respectful of the expected mother's privacy.
- Assess the patient for peripheral edema. This may indicate Pre-eclampsia / Eclampsia. Monitor patient closely and watch for seizure activity.
- Tag the mother and baby with the same information. Consider keeping a zip lock bag with name tags/ Sharpie attached to the O.B. kit itself.
- Green or brown amniotic fluid indicates the presence of Meconium (fetal stool) and should be reported immediately to the receiving facility staff.

## Obstetrical Complications Protocol

Obstetrical complications can rapidly lead to hypovolemic shock and threaten the life of the mother and child. Care should be focused on assessing the situation, initiating routine patient care, and beginning treatment for shock. Monitor vitals closely.

### Placenta Previa

*Placenta previa* occurs because of abnormal implantation of the placenta on the lower half of the uterine wall. Bleeding occurs when the lower uterus begins to contract and dilate in preparation for labor and pulls the placenta away from the uterine wall. The hallmark of *placenta previa* is the onset of painless bright red vaginal bleeding, usually in the 3<sup>rd</sup> trimester of pregnancy.

### EMR Care

1. **Initiate** Paramedic Care intercept.
2. **Oxygen:** Titrate O2 admin to maintain  $SpO_2 \geq 94\%$ .
3. Note the amount of bleeding.
4. Place the patient on her **left side**.
5. Load and transport as soon as possible.

### EMT Care

1. EMT Care includes all components of *EMR Care*
2. **Contact Medical Control** as soon as possible.

### A-EMT/ EMT-I Care

1. A-EMT/ EMT-I includes all components of *EMT Care*.
2. **IV Fluid Therapy:** 500mL fluid boluses to maintain a systolic BP of at least 100mmHg.

### Paramedic Care

1. Paramedic Care includes all components of *A-EMT/ EMT-I Care*.



Memorial EMS  
Decatur Memorial EMS  
Springfield Memorial EMS

## Obstetrical Complications Protocol

### Ectopic Pregnancy

*Ectopic Pregnancy* refers to the abnormal implantation of the fertilized egg outside of the uterus, usually in the fallopian tube. It can be a life-threatening condition and accounts for approximately 10% of maternal mortality in the 1<sup>st</sup> trimester. Typical presentation occurs in weeks 4-11 after LMP. Patient may not know she is pregnant.

Ectopic pregnancy presents as abdominal pain which starts out as diffuse tenderness and then localizes as a sharp pain in the lower abdomen on the effected side. Assume that any female of childbearing age with lower abdominal pain is experiencing an ectopic pregnancy.

### EMR Care

1. **Oxygen:** Titrate O2 admin to maintain SpO2  $\geq$  94%.
2. **Initiate** Paramedic Care intercept.

### EMT Care


1. EMT Care includes all components of *EMR Care*.
2. **Contact Medical Control** as soon as possible.

### A-EMT/ EMT-I Care

3. A-EMT/ EMT-I includes all components of *EMT Care*.
4. **IV Fluid Therapy:** 500mL fluid boluses to maintain a systolic BP of at least 100mmHg.

### Paramedic Care

1. Paramedic Care includes all components of *A-EMT/ EMT Care*.



Memorial EMS  
Decatur Memorial EMS  
Springfield Memorial EMS  
**Obstetrical Complications Protocol**

## **Abruptio Placentae**

*Abruptio placentae* is the premature separation of a normally implanted placenta from the uterine wall. Signs and symptoms can vary depending on the extent and character of the abruption. Abruptio placentae can be caused by even minor trauma. Bleeding can be massive.

### **EMR Care**

1. **Oxygen:** Titrate O<sub>2</sub> admin to maintain SpO<sub>2</sub>  $\geq$  94%.
2. Note the amount of bleeding.
3. Place the patient on her left side.
4. Load and transport as soon as possible.
5. **Initiate** Paramedic Care intercept.

### **EMT Care**

1. EMT Care includes all components of *EMR Care*.
2. **Contact Medical Control** as soon as possible.

### **A-EMT/ EMT-I Care**

1. A-EMT/ EMT-I includes all components of *EMT Care*.
2. **IV Fluid Therapy:** 500mL fluid boluses to maintain a systolic BP of at least 100mmHg.
3. Establish a 2<sup>nd</sup> IV enroute if time permits.

### **Paramedic Care**

1. Paramedic Care includes all components of *A-EMT/ EMT-I Care*.

## **Antepartum & Postpartum Hypertension**

*Pre-eclampsia* is defined as an increase in systolic blood pressure by 30mmHg and/or a diastolic increase of 15mmHg over baseline on at least two occasions at least 6 hours apart. *Pre-eclampsia* is most seen in the last 10 weeks of gestation and is thought to be caused by abnormal vasospasm.

*Pre-Eclampsia:* Characterized by hypertension and edema to the hands and face (and protein in the urine).

*Severe Pre-Eclampsia:* Characterized by marked hypertension (160/100 or higher), generalized edema, headache, visual disturbances, pulmonary edema, and a dramatic decrease in urine output (along with a significant increase of protein in the urine). May also present with RUQ pain.

*Eclampsia:* Characterized by generalized tonic-clonic seizure activity often preceded by flashing lights or spots before the eyes. Altered Mental Status may be present. Can occur anytime during pregnancy and up to six weeks AFTER delivery.

## Obstetrical Complications Protocol

### Antepartum & Postpartum Hypertension {Continued}

Antepartum Hypertension: Characterized by pregnancy with SBP > 140 or DPB > 90, headache, visual complications, AMS, stroke symptoms, or seizures.

Postpartum Hypertension: Characterized by pregnancy with SBP > 140 or DPB > 90, headache, visual complications, AMS, stroke symptoms, or seizures in the post-delivery patient for up to six weeks.

**Note:** The risk of fetal mortality increases by 10% with each maternal seizure.

### EMR Care

1. Assure minimal CNS stimulation to prevent seizures (*i.e.*, do not check papillary light reflex).
2. Place the patient on her left side (if pregnant).
3. Load and transport as soon as possible.
4. **Initiate** Paramedic Care intercept.

### EMT Care

1. EMT Care includes all components of *EMR Care*.
2. **Contact Medical Control** as soon as possible.

### A-EMT/ EMT-I Care

1. A-EMT/ EMT-I includes all components of *EMT Care*.
2. **IV Fluid Therapy:** TKO.
3. If the patient is actively seizing, refer to the Seizure Protocol.

### Paramedic Care

1. Paramedic Care includes all components of *A-EMT/ EMT-I Care*
2. **Contact Medical Control** for possible **Magnesium Sulfate** administration orders.

### Transporting Units

Hypertensive antepartum and postpartum patients are especially challenging patients who can have very complex care needs. As such, transport to the most appropriate facility is necessary to provide the patient the best treatment options.

1. Patients meeting *Patient Destination Criteria (6.G.1)* should be transported to a Level III Perinatal facility if possible.
  - a. If unable to safely transport directly, include this in PCR documentation.
  - b. If the patient refuses, include documentation of the informed refusal in communication with **Medical Control** and in the PCR.

## Abnormal Delivery Protocol

Abnormal delivery situations can be especially challenging in the pre-hospital setting. Care should be focused on initiating *Routine Patient Care* to treat for shock and rapid transport to the hospital.

### Breech Presentation

A *breech* presentation is the term used to describe a situation in which either the buttocks or both feet present first.

### EMR Care

1. Render initial care in accordance with the *Routine Patient Care Protocol*.
2. **Oxygen:** If respiratory distress is noted, 15 LPM via NRM or if unable to tolerate the mask, 6 LPM via nasal cannula.
  - a. If no obvious respiratory distress is noted, apply a pulse ox. If  $\geq 94\%$  and no signs/ symptoms of respiratory distress, no Oxygen is required. If  $\leq 94\%$  apply nasal cannula at 2-6 LPM or 15 LPM via NRM as needed to raise pulse ox to  $\geq 94\%$ .
3. Load and transport as soon as possible.
4. **Initiate** Paramedic Care intercept.
5. Never attempt to pull the baby from the vagina by the trunk or legs.
6. As soon as the legs are delivered, support the baby's body (wrapped in a towel).
7. After the shoulders are delivered, gently elevate the trunk and legs to aid in the delivery of the head.
8. The head should deliver in 30 seconds. If it does not – reach 2 fingers into the vagina to locate the infant's mouth. Press the vaginal wall away from the baby's mouth to provide unrestricted respirations.
9. **Contact Medical Control** as soon as possible.

### EMT Care

1. EMT Care includes all components of *EMR Care*.

### A-EMT/ EMT-I Care

1. A-EMT/ EMT-I includes all components of *EMT Care*.

### Paramedic Care

1. Paramedic Care includes all components of *A-EMT/ EMT-I Care*

### Shoulder Dystocia

Shoulder dystocia is a halting of the natural progress of delivery due to failure to deliver the baby's shoulders. This occurs when the anterior shoulder becomes stuck on the mother's pubic symphysis. Occurs in up to 3% of deliveries. Failure to deliver the anterior shoulder in a timely fashion can result in permanent brachial plexus injury, fetal hypoxia, and death. Risk factors for dystocia are a large for gestational age infant (estimated weight >3500 grams), maternal diabetes and maternal obesity.

#### Signs you should be concerned about shoulder dystocia:

- Turtle sign: delivery of the fetal head followed by retraction of the head into the vaginal canal
- >60 seconds between delivery of the fetal head and delivery of the shoulders

Memorial EMS  
Decatur Memorial EMS  
Springfield Memorial EMS  
**Abnormal Delivery Protocol**

**Shoulder Dystocia (cont.)**

**EMR Care**

1. Render initial care in accordance with the *Routine Patient Care Protocol*.
2. **Oxygen:** If respiratory distress is noted, 15 LPM via NRM or if unable to tolerate the mask, 6 LPM via nasal cannula.
  - a. If no obvious respiratory distress is noted, apply a pulse ox. If  $\geq 94\%$  and no signs/ symptoms of respiratory distress, no Oxygen is required. If  $\leq 94\%$  apply nasal cannula at 2-6 LPM or 15 LPM via NRM as needed to raise pulse ox to  $\geq 94\%$ .
3. **Initiate** Paramedic Care intercept. **If you are concerned about the possibility of shoulder dystocia update medical control of a complicated delivery.**

**EMT Care**

1. **If you are concerned about the possibility of shoulder dystocia update medical control of a complicated delivery.**
2. **Contact Medical Control** as soon as possible.

**A-EMT/ EMT-I Care**

1. A-EMT/ EMT-I includes all components of *EMT Care*.

**Paramedic Care**

1. Paramedic Care includes all components of *A-EMT/ EMT-I Care*

**Steps to assist in shoulder delivery:**

**McRoberts Maneuver**— hyperflexion of hips creates superior displacement of the pubic symphysis and sacral extension. Additional gentle downward suprapubic pressure helps further disengage the stuck anterior shoulder.

1. Ask mother to pull her knees to her chest keeping her thighs against her abdomen.
2. Make a fist and apply gentle downward suprapubic (not fundal) pressure to help disengage the stuck shoulder.
3. If these maneuvers fail to assist in delivery move on to the next maneuver

**Rubin II Maneuver** – gentle sweeping of the anterior shoulder in the direction the baby is facing to disengage the anterior shoulder from the pubic symphysis.

1. Insert two fingers into the vaginal canal to the between baby's head and maternal pubic symphysis.
2. Gently push the shoulder in the direction the baby is facing to disengage the shoulder from the symphysis.

Memorial EMS  
Decatur Memorial EMS  
Springfield Memorial EMS  
**Abnormal Delivery Protocol**

### **Prolapsed Cord**

A *prolapsed cord* occurs when the umbilical cord precedes the fetal presenting part. This causes the cord to be compressed between the fetus and the pelvis and blocks fetal circulation. Fetal death will occur quickly without prompt intervention.

### **EMR Care**

1. Render initial care in accordance with the *Routine Patient Care Protocol*.
2. **Oxygen:** If respiratory distress is noted, 15 LPM via NRM or if unable to tolerate the mask, 6 LPM via nasal cannula.
  - a. If no obvious respiratory distress is noted, apply a pulse ox. If  $\geq 94\%$  and no signs/ symptoms of respiratory distress, no Oxygen is required. If  $\leq 94\%$  apply nasal cannula at 2-6 LPM or 15 LPM via NRM as needed to raise pulse ox to  $\geq 94\%$ .
3. **Initiate** Paramedic Care intercept.
4. Place the mother in Trendelenburg Position.
5. **Do not pull on the cord and do not attempt to push the cord back into the vagina.**
6. Place a gloved finger/hand in the vagina between the pubic bone and the presenting part with the cord between the fingers and exert counter pressure against the presenting part.
7. Palpate the cord for pulsations.
8. Keep the exposed cord warm and moist.
9. Keep the hand in position and transport immediately.

### **EMT Care**

1. EMT Care includes all components of *EMR Care*.
2. **Contact Medical Control** as soon as possible.

### **A-EMT/ EMT-I Care**

1. A-EMT/ EMT-I includes all components of *EMT Care*.

### **Paramedic Care**

1. Paramedic Care includes all components of *A-EMT/ EMT-I Care*

## Abnormal Delivery Protocol

### Limb Presentation

Although relatively uncommon, the baby may be lying transverse across the uterus. In these cases, an arm or leg is the presenting part protruding from the vagina and will require delivery by cesarean section. **Under no circumstances should you attempt a field delivery** with a limb presentation.

### EMR Care

1. Render initial care in accordance with the *Routine Patient Care Protocol*.
2. **Oxygen:** If respiratory distress is noted, 15 LPM via NRM or if unable to tolerate the mask, 6 LPM via nasal cannula.
  - a. If no obvious respiratory distress is noted, apply a pulse ox. If  $\geq 94\%$  and no signs/ symptoms of respiratory distress, no Oxygen is required. If  $\leq 94\%$  apply nasal cannula at 2-6 LPM or 15 LPM via NRM as needed to raise pulse ox to  $\geq 94\%$ .
3. **Initiate** Paramedic Care intercept.
4. Place the mother in the Trendelenburg Position.
5. Avoid touching the limb (doing so may stimulate the infant to gasp). **Do not pull on the extremity and do not attempt to push the limb back into the vagina.**

### EMT Care

1. EMT Care includes all components of *EMR Care*.
2. **Contact Medical Control** as soon as possible.

### A-EMT/ EMT-I Care

1. A-EMT/ EMT-I includes all components of *EMT Care*.

### Paramedic Care

1. Paramedic Care includes all components of *A-EMT/ EMT-I Care*